

Technical Data Sheet

Biotinylated Human IL13RA2 Protein (C-Fc-Avi)

Catalog Number: 804803, 804804

Size: 25 ug, 100 ug

Target Name: IL13RA2, CD213A2, IL-13R, IL13BP

Regulatory Status: RUO

Product Details

Application: ELISA, BLI

Format: Liquid, Biotinylated

Expression Host: CHO

Species: Human

Sources: Recombinant Human IL13RA2 protein (Cys22-Leu342) with C-terminus Fc-Avi tag is expressed in CHO cells. This protein was site-specifically labeled with Biotin by BirA ligase.

Accession Number: Q14627

Molecular Weight: The protein has a predicted molecular weight of 65.7 kDa. Under DTT-reducing conditions, it migrates at approximately 65-80 kDa on SDS-PAGE.

Affinity Tag: C-Fc-Avi

Purity: >95% based on SDS-PAGE under reducing condition

Formulation: 1xPBS buffer, pH7.4, 0.22 µm filtered

Endotoxin level: Not tested

Protein Concentration: 25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.


Storage and Handling: Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

Background Information

CD213A2, also known as IL13R?2, is a type I transmembrane protein belonging to the hematopoietin receptor family. It binds interleukin-13 (IL-13) with high affinity but lacks a functional cytoplasmic signaling domain, suggesting that it primarily acts as a decoy receptor, antagonizing IL-13 signaling mediated by the IL-13R?1/IL-4R? complex. IL13R?2 is expressed in fibroblasts, smooth muscle cells, keratinocytes, and activated B cells, though its surface expression is tightly regulated and much of it resides intracellularly or in soluble form. In addition to inhibiting IL-13 activity, IL13R?2 has been reported to suppress IL-4 signaling via physical interaction with IL-4R?, while paradoxically promoting TGF-? production and fibrosis. Importantly, IL13R?2 is highly and specifically overexpressed in certain cancers, such as glioblastoma multiforme and high-grade astrocytomas, making it a promising target for tumor-specific immunotherapies and viral vector-mediated gene delivery.

approaches.

Product Data



Human Human IL13RA2 Protein (C-Fc-Avi) Protein is biotinylated by BirA ligase in vitro. Biotinylated Human Human IL13RA2 protein on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The purity of this protein appears to be greater than 95%. Based on Gel shift Assay by co-incubation with Streptavidin, biotinylation efficiency is >70% for Biotinylated Human Human IL13RA2.