

## Technical Data Sheet

### PB Anti-Human CD127

**Catalog Number:** 104803, 104804

**Size:** 25 tests, 100 tests

**Target Name:** CD127, IL-7 receptor  $\alpha$  chain, IL-7R $\alpha$

**Regulatory Status:** RUO

#### Product Details

---

**Clone:** 127AB

**Application:** FC

**Reactivity:** Human

**Format:** PB

**Isotype:** Rabbit IgG

**Antibody Type:** Monoclonal

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA

**Protein Concentration:** Supplied at a lot-specific concentration.

**Storage and Handling:** The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.

**Recommended Usage:** For flow cytometric staining, it is recommended to use 5  $\mu$ L of this reagent per 0.5-1.0 million cells in a 100  $\mu$ L volume. Optimal reagent performance should be determined by titration for each specific application.

**Excitation Laser:** Violet laser (405nm)

#### Background Information

---

The interleukin-7 receptor is a protein found on the surface of cells. It is made up of two different smaller protein chains - i.e. it is a heterodimer, and consists of two subunits, interleukin-7 receptor- $\alpha$  (CD127) and common- $\gamma$  chain receptor (CD132). The common- $\gamma$  chain receptors is shared with various cytokines, including interleukin-2, -4, -9, and -15. Interleukin-7 receptor is expressed on various cell types, including naive and memory T cells and many others. Interleukin-7 receptor has been shown to play a critical role in the development of immune cells called lymphocytes.