

## InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

## **Technical Data Sheet**

PE conjugated Human CD32a R167 /Fc gamma RIIa (C-His)

Catalog Number: 820401, 820402

Size: 25 ug, 100 ug

Target Name: CD32a, FCGR2A, CD32, FCG2, FCGR2A1, IGFR2

Regulatory Status: RUO

## **Product Details**

Application: FC
Format: Liquid, PE
Expression Host: CHO

Species: Human

Sources: Recombinant CD32a R167 /Fc gamma RIIa (Ala 36 - Ile 218) with C-terminus His-tag is

expressed in CHO cell and conjugated to PE.

**Accession Number: Q92835** 

Molecular Weight: The protein has a predicted molecular weight of 21.9 kDa. Under DTT-reducing

conditions, it migrates at approximately 35 kDa on SDS-PAGE prior to conjugation.

Affinity Tag: C-His

Formulation: 1xPBS buffer, pH7.4, 0.09% NaN3 with a carrier protein

Endotoxin level: Not tested

Protein Concentration: 25μg size is bottled at 0.1mg/mL concentration. 100 μg size is bottled at lot

specific concentration.

Storage and Handling: Briefly centrifuge the vial upon receipt. An unopened vial may be stored at

2-8°C for up to six months.

## **Background Information**

Fc?RII, or CD32, is a low-affinity receptor for IgG Fc region. In human, it is expressed as three isoforms (A, B, C). The activating isoform, Fc?RIIA (CD32a), is found on monocytes, macrophages, granulocytes, platelets, and mast cells. It features an extracellular domain, transmembrane region, and a cytoplasmic tail with a non-classical ITAM, crucial for dimerization and signaling. Fc?RIIA plays a key role in inflammation, phagocytosis, and platelet activation, influencing conditions like thrombocytopenia, rheumatoid arthritis, and lupus. Its activity is modulated by lipid rafts, and the receptor's affinity for IgG subclasses varies.