www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

Human CD32b/c (FcγRIIB) Protein (C-His-Avi)

Catalog Number:	820501, 820502
Size:	25 ug, 100 ug
Target Name:	CD32b/c, FCGR2B, C, FcRII-b, c, CD32, FCG2, IGFR2
Regulatory Status:	RUO

PRODUCT DETAILS

Application:	ELISA, BLI
Format:	Liquid, Purified
Expression Host:	CHO
Species:	Human
Sources:	Recombinant CD32b/c /Fc gamma RIIb (Ala46-Pro217) with C-terminus Avi-His-tag is expressed in CHO cell.
Accession Number:	P31994
Molecular Weight:	The protein has a predicted molecular weight of 23.5 kDa. Under DTT-reducing conditions, it migrates at approximately 35 kDa on SDS-PAGE.
Affinity Tag:	C-Avi-His
Purity:	>95% based on SDS-PAGE under reducing condition
Formulation:	1xPBS buffer, pH7.4, 0.22 μm filtered
Endotoxin level:	Not tested
Protein Concentration:	25μg size is bottled at 0.2mg/mL concentration. 100 μg size is supplied at a lot-specific concentration.
Storage and Handling:	Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 μm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ –70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

BACKGROUND INFORMATION

CD32B (FCGR2B) and CD32C (FCGR2C) are 40 kDa, type I transmembrane proteins belonging to the Ig superfamily of low-affinity IgG Fc receptors. CD32B contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic tail, while CD32C has an immunoreceptor tyrosine-based activation motif (ITAM). Both receptors can bind monomeric IgG and IgG complexes, and are expressed on B cells, monocytes, macrophages, granulocytes, platelets, and mast cells. CD32B primarily acts as a negative regulator, inhibiting immune cell activation, proliferation, endocytosis, phagocytosis, and degranulation. In contrast, CD32C has an activating function, contributing to immune responses. Together, they play crucial roles in regulating immune cell functions and

maintaining immune homeostasis. CD32B and CD32C share almost identical extracellular domain. The aa sequence of Fc gamma RIIB and Fc gamma RIIC are the same for the region expressed (Ala46-Pro217)

PRODUCT DATA



Human CD32b/c /Fc gamma RIIb (C-His-Avi) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The purity of this protein appears to be greater than 95%.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.