

Technical Data Sheet

PE conjugated Human CD32b/c /Fc gamma RIIB (C-His)

Catalog Number: 820701, 820702

Size: 25 ug, 100 ug

Target Name: CD32b/c, FCGR2B, C, FcRII-b, c, CD32, FCG2, IGFR7

Regulatory Status: RUO

Product Details

Application: FC

Format: Liquid, PE

Expression Host: CHO

Species: Human

Sources: Recombinant CD32b/c /Fc gamma RIIB (Ala46-Pro217) with C-terminus His-tag is expressed in CHO cell and conjugated to PE.

Accession Number: P31994

Molecular Weight: The protein has a predicted molecular weight of 20.9 kDa. Under DTT-reducing conditions, it migrates at approximately 35 kDa on SDS-PAGE prior to conjugation.

Affinity Tag: C-His

Formulation: 1xPBS buffer, pH7.4, 0.09% NaN₃ with a carrier protein

Endotoxin level: Not tested

Protein Concentration: 25µg size is bottled at 0.1mg/mL concentration. 100 µg size is bottled at lot specific concentration.

Storage and Handling: Briefly centrifuge the vial upon receipt. An unopened vial may be stored at 2–8°C for up to six months.

Background Information

CD32B (FCGR2B) and CD32C (FCGR2C) are 40 kDa, type I transmembrane proteins belonging to the Ig superfamily of low-affinity IgG Fc receptors. CD32B contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic tail, while CD32C has an immunoreceptor tyrosine-based activation motif (ITAM). Both receptors can bind monomeric IgG and IgG complexes, and are expressed on B cells, monocytes, macrophages, granulocytes, platelets, and mast cells. CD32B primarily acts as a negative regulator, inhibiting immune cell activation, proliferation, endocytosis, phagocytosis, and degranulation. In contrast, CD32C has an activating function, contributing to immune responses. Together, they play crucial roles in regulating immune cell functions and maintaining immune homeostasis. CD32B and CD32C share almost identical extracellular domain. The aa sequence of Fc gamma RIIB and Fc gamma RIIC are the same for the region expressed (Ala46-Pro217)