

Technical Data Sheet

Cynomolgus Monkey CD200 Protein (C-Fc)

Catalog Number: 605601, 605602
Size: 25 ug, 100 ug
Target Name: CD200, MOX1, MOX2, MRC, OX-2, My033
Regulatory Status: RUO

Product Details


Application: ELISA, BLI
Format: Liquid, Purified
Expression Host: CHO
Target Name: CD200, MOX1, MOX2, MRC, OX-2, My033
Species: Cynomolgus monkey
Accession Number: □EHH51019
Sources: Recombinant Cynomolgus Monkey CD200 (Gln56-Gly257) with C-terminus Fc tag is expressed in CHO cells.
Molecular Weight: This protein has a predicted molecular weight of 48.8 kDa. Under DTT-reducing conditions, the protein migrates at approximately 65 kDa on SDS-PAGE.
Affinity Tag: C-Fc
Purity: >95% based on SDS-PAGE under reducing condition
Regulatory Status: RUO
Formulation: 1xPBS buffer, pH7.4, 0.22 µm filtered
Endotoxin level: Less than 0.1 EU/µg protein as determined by the LAL method
Protein Concentration: 25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.
Storage and Handling: Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

Background Information


CD200 (OX-2) is a cell surface glycoprotein that regulates immune responses through its receptor, CD200R, mainly expressed on myeloid cells. It plays a crucial role in suppressing alloimmune and autoimmune responses, contributing to immune privilege in various tissues. CD200-CD200R signaling is essential in regulating anti-tumor immunity, with overexpression linked to malignancies like chronic lymphocytic leukemia (CLL) and cancer stem cells. Additionally, CD200 signaling is involved in the central nervous system, particularly in diseases like Parkinson's, where it affects microglia activation. Elevated CD200 expression is associated with reduced transplant rejection, autoimmunity, and allergic diseases, but may also promote tumor cell survival. While it

helps prevent graft rejection and autoimmune diseases, high CD200 levels in cancers have been linked to poor prognosis. This dual role underscores the complexity of CD200 in immune regulation and its potential as both a therapeutic target and a biomarker.

Product Data



Purified Cynomolgus Monkey CD200 Protein (C-Fc) on SDS-PAGE under reducing (P+) and non-reducing (P-) conditions. The purity of the purified protein appears to be greater than 95% based on reducing condition.



Human CD200R (C-His) is coated at 2 ug/mL (200 ng/well). Cyno Monkey CD200 (C-Fc) can bind human CD200R in a dose-dependent manner with the ED50 of 2-20 ng/mL