

Technical Data Sheet

Human Notum Protein (C-Fc)

Catalog Number: 606001, 606002

Size: 25 ug, 100 ug

Target Name: Notum, Palmitoleoyl-protein carboxylesterase NOTUM

Regulatory Status: RUO

Product Details

Application: ELISA, BLI

Format: Liquid, Purified

Expression Host: CHO

Species: Human

Accession Number: Q6P988

Sources: Recombinant Human Notum (Ser20-Ser496) with C-terminus Fc tag was expressed in CHO cells.

Molecular Weight: This protein has the predicted molecular weight of 79.89 kD. Under DTT-reducing conditions, the protein migrates at approximately 80 kD on SDS-PAGE

Affinity Tag: C-Fc

Purity: >80% based on SDS-PAGE under reducing condition

Formulation: 1xPBS buffer, pH7.4, 0.22 µm filtered

Endotoxin level: Not tested


Protein Concentration: 25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.

Storage and Handling: Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

Background Information

Notum is a secreted carboxylesterase enzyme that acts as a negative regulator of the Wnt signaling pathway. It plays a crucial role in development, tissue homeostasis, and regeneration by modulating Wnt activity. Notum inhibits Wnt signaling by removing an essential lipid modification (palmitoleate) from Wnt proteins, in this manner preventing their binding to receptors and subsequent activation of downstream pathways. Dysregulation of Notum has been linked to various diseases, including cancer and degenerative disorders, making it a promising target for therapeutic intervention..

Product Data



Purified Human Notum Protein (C-Fc) on SDS-PAGE under reducing (P+) and non-reducing (P-) conditions. The purity of the purified protein appears to be greater than 80% based on reducing condition.