

## InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

## **Technical Data Sheet**

Human GITRL/TNFSF18 (N-His-Avi)

Catalog Number: 606601, 606602

Size: 25 ug, 100 ug

Target Name: TNFSF18, GITR Ligand, AITRL, TL6, GITRL

Regulatory Status: RUO

#### **Product Details**

Application: ELISA, BLI Format: Liquid, Purified Expression Host: CHO

Species: Human

Accession Number: Q9UNG2

Sources: Recombinant Human TNFSF18 (Glu52-Ser177) with N-terminus His-Avi tag is expressed

in CHO cells.

Molecular Weight: This prtoein has the predicted molecular weight of 18.3 kD. Under DTT-reducing

conditions, the protein migrates at approximately 20 kD on SDS-PAGE

**Affinity Tag:** N-His-Avi

Purity: >95% based on SDS-PAGE under reducing condition

Formulation: 1xPBS buffer, pH7.4, 0.22 µm filtered

Endotoxin level: Less than 0.1 EU/µg protein as determined by the LAL method

Protein Concentration: 25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a

lot-specific concentration.

**Storage and Handling:** Briefly centrifuge the vial upon receipt. An unopened vial can be stored at  $4^{\circ}$ C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22  $\mu$ m-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at <=  $-70^{\circ}$ C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

#### **Background Information**


Human GITRL (Glucocorticoid-Induced TNFR-Related Ligand), also known as TNFSF18, is a member of the tumor necrosis factor (TNF) superfamily. It is a type II transmembrane protein primarily expressed on activated antigen-presenting cells, including B cells, dendritic cells, and macrophages. GITRL binds to its receptor GITR, which is found on activated T cells, regulatory T cells (Tregs), and natural killer (NK) cells. The GITR–GITRL interaction delivers a co-stimulatory signal that promotes T cell activation, proliferation, and survival, while also modulating the suppressive activity of Tregs. This signaling pathway plays a key role in immune regulation, inflammation, and antitumor immunity.



# InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

### **Product Data**



Human GITRL (TNFSF18) (N-His-Avi) on SDS-PAGE under reducing condition (P+) and non-reducing condition (P-). The gel was stained for 1 hour with BlinkBlue (catalog 700102). The purity of this protein appears to be greater than 95%.