

## InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

### **Technical Data Sheet**

APC Anti-Mouse CD3

Catalog Number: 200113, 200114

Size: 25 tests, 100 tests

Target Name: CD3, T cell antigen receptor complex, T3

Regulatory Status: RUO

#### **Product Details**

Clone: KT3
Application: FC
Reactivity: Mouse
Format: APC
Isotype: Rat IgG2a

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v)

BSA

**Protein Concentration:** Supplied at a lot-specific concentration.

Storage&Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

**Recommended Usage:** For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by

titration for each specific application. **Excitation Laser:** Red Laser (633 nm)

### **Background Information**


CD3 (cluster of differentiation 3) is a protein complex and T cell co-receptor that is involved in activating both the cytotoxic T cell (CD8+ naive T cells) and T helper cells (CD4+ naive T cells). CD3 is a member of the Ig superfamily and primarily expressed on T cells, NK-T cells, and at different levels on thymocytes during T cell differentiation. CD3 is composed of CD3?, ?, ? and ? chains. It forms a TCR complex by associating with TCR ?/? or ?/? chains. CD3 plays a critical role in TCR signal transduction, T cell activation, and antigen recognition by binding the peptide/MHC antigen complex

#### **Product Data**



# InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Mouse splenocytes stained with either APC Anti-Mouse CD3 clone KT3 (purple histogram) or an isotype control (gray histogram).