

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

Technical Data Sheet

iF488 Anti-Human CD127

Catalog Number: 104807, 104808

Size: 25 tests, 100 tests

Target Name: CD127, IL-7 receptor ? chain, IL-7R?

Regulatory Status: RUO

Product Details

Clone: 127AB
Application: FC
Reactivity: Human
Format: iF488
Isotype: Rabbit IgG

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v)

BSA

Protein Concentration: Supplied at a lot-specific concentration.

Storage&Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

Recommended Usage: For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 μ L volume. Optimal reagent performance should be determined by

titration for each specific application. **Excitation Laser:** Blue Laser (488 nm)

Background Information


The interleukin-7 receptor is a protein found on the surface of cells. It is made up of two different smaller protein chains - i.e. it is a heterodimer, and consists of two subunits, interleukin-7 receptor-? (CD127) and common-? chain receptor (CD132). The common-? chain receptors is shared with various cytokines, including interleukin-2, -4, -9, and -15. Interleukin-7 receptor is expressed on various cell types, including naive and memory T cells and many others. Interleukin-7 receptor has been shown to play a critical role in the development of immune cells called lymphocytes.

Product Data



InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Human peripheral blood lymphocytes stained either iF488 Anti-Human CD127 clone 127AB (color-filled histogram) or an isotype control (gray histogram).