

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

Technical Data Sheet

APC Anti-Human CD64

Catalog Number: 109311, 109312

Size: 25 tests, 100 tests

Target Name: CD64, FCGR1A, FCG1, FCGR1, IGFR1

Regulatory Status: RUO

Product Details

Clone: H22 Application: FC Reactivity: Human Format: APC

Isotype: Mouse IgG1

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v)

BSA

Protein Concentration: Supplied at a lot-specific concentration.

Storage&Handling: The antibody solution should be stored undiluted between 2°C and 8°C, and

protected from prolonged exposure to light. Do not freeze.

Recommended Usage: For flow cytometric staining, it is recommended to use 5 uL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by

titration for each specific application. **Excitation Laser:** Red Laser (633 nm)

Background Information


CD64, also known as Fc?RI or FcR I, is a 72 kDa type I glycoprotein and a member of the immunoglobulin superfamily. This high-affinity IgG Fc receptor is predominantly expressed on monocytes, macrophages, dendritic cells, and activated granulocytes. Its expression can be upregulated by IFN-? stimulation, enhancing its role in immune responses. CD64 binds IgG immune complexes and is involved in several crucial immune functions, including antigen capture, phagocytosis of IgG/antigen complexes, and antibody-dependent cellular cytotoxicity (ADCC). By mediating these processes, CD64 contributes to the activation of innate immune responses and the clearance of immune complexes.

Product Data



InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Human peripheral blood monocytes stained either APC Anti-Human CD64 clone H22 (color-filled histogram) or an isotype control (gray histogram).