

## Anti-human/mouse/rat CD278 (ICOS) Antibody

<b>Catalog Number:</b>	111201, 111202
<b>Size:</b>	25 ug, 100 ug
<b>Target Name:</b>	CD278, ICOS, Inducible Costimulatory molecule
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Clone:</b>	C398.4A
<b>Application:</b>	Flow Cytometry
<b>Reactivity:</b>	Human, mouse, rat
<b>Format:</b>	Purified
<b>Isotype:</b>	Armenian Hamster IgG
<b>Antibody Type:</b>	Monoclonal
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide
<b>Protein Concentration:</b>	0.5 mg/mL
<b>Storage&amp;Handling:</b>	The antibody solution should be stored between 2°C and 8°C
<b>Recommended Usage:</b>	For flow cytometric staining, it is recommended to use less than 0.2 ug of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application.
<b>RRID:</b>	AB_3738962

### BACKGROUND INFORMATION

---

ICOS is a 47-57 kDa costimulatory molecule homologous to CD28/CTLA-4 and is expressed on activated T cells and a subset of thymocytes. Engagement of ICOS enhances T cell proliferation, promotes cytokine production such as IL-4 and IL-10, and plays an important role in humoral immunity and Th2 cell differentiation.