

## In Vivo Star Anti-Mouse CD152 (CTLA-4) Antibody

|                           |                        |
|---------------------------|------------------------|
| <b>Catalog Number:</b>    | 509901, 509902, 509903 |
| <b>Size:</b>              | 1 mg, 5 mg, 25 mg      |
| <b>Target Name:</b>       | mouse CTLA-4, CD152    |
| <b>Regulatory Status:</b> | RUO                    |

### PRODUCT DETAILS

---

|                             |   |
|-----------------------------|---|
| <b>Clone:</b>               | 9D9   |
| <b>Application:</b>         | ELISA, WB, Flow cytometry, IHC, ICC, animal model study   |
| <b>Reactivity:</b>          | Mouse   |
| <b>Format:</b>              | Liquid  |
| <b>Product Description:</b> | In vivo Grade Recombinant Anti-mouse CTLA-4 Monoclonal Antibody   |
| <b>Isotype:</b>             | Mouse IgG2b Kappa   |
| <b>Antibody Type:</b>       | Recombinant   |
| <b>Purity:</b>              | >95% by reducing SDS-PAGE   |
| <b>Endotoxin:</b>           | < 1 EU per 1 mg of the protein by the LAL method.   |
| <b>Storage Conditions:</b>  | 4°C   |
| <b>Grade:</b>               | In vivo   |
| <b>Recommended Usage:</b>   | This product is suitable for in vivo animal use. Optimal amounts need to be determined empirically for each experiment. |
| <b>Hidden Synonyms:</b>     | InVivoMab, InVivoPlus, GoInVivo, In Vivo Gold   |
| <b>RRID:</b>                | AB_3739373  |

### BACKGROUND INFORMATION

---

Cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), also known as CD152, is a critical immune checkpoint receptor that functions as a negative regulator of T cell activation. It is primarily expressed on activated CD4+ and CD8+ T cells and is constitutively expressed at high levels on regulatory T cells (Tregs). CTLA-4 plays a central role in maintaining immune homeostasis by limiting excessive T cell responses and promoting peripheral tolerance.

Structurally, CTLA-4 is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily. Its extracellular region consists of a single IgV-like domain responsible for ligand binding, followed by a transmembrane region and a short cytoplasmic tail. The cytoplasmic domain lacks intrinsic enzymatic activity but contains conserved signaling motifs, including a tyrosine-based motif that mediates interactions with intracellular signaling and trafficking proteins. CTLA-4 predominantly resides in intracellular vesicles and is rapidly transported to the cell surface following T cell activation.

The primary ligands for CTLA-4 are the B7 family co-stimulatory molecules CD80 (B7-1) and CD86 (B7-2), which are expressed on

antigen-presenting cells such as dendritic cells, macrophages, and B cells. CTLA-4 binds CD80 and CD86 with significantly higher affinity and avidity than the activating receptor CD28. By outcompeting CD28 for ligand binding and actively removing CD80/CD86 from the surface of antigen-presenting cells through trans-endocytosis, CTLA-4 effectively dampens co-stimulatory signaling and restrains T cell activation.

Dysregulation of CTLA-4 function is associated with a range of diseases. Genetic deficiency or loss-of-function mutations in CTLA-4 can lead to severe lymphoproliferative disorders, autoimmunity, and immune dysregulation due to uncontrolled T cell activation. Conversely, excessive CTLA-4 activity can contribute to impaired immune responses, including reduced anti-tumor immunity. In cancer, tumor-induced upregulation of CTLA-4 signaling contributes to immune evasion by suppressing effective T cell responses.

CTLA-4 is a landmark target in immunotherapy. Therapeutic antibodies that block CTLA-4, such as immune checkpoint inhibitors, enhance T cell activation and proliferation by restoring co-stimulatory signaling, leading to improved anti-tumor immune responses in several cancers. However, CTLA-4 blockade can also disrupt immune tolerance, resulting in immune-related adverse events. Conversely, strategies that enhance CTLA-4 function or signaling are being explored for the treatment of autoimmune and inflammatory diseases. Together, these approaches highlight CTLA-4's pivotal role at the intersection of immune regulation, disease, and therapy.

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.