

## FITC Anti-Mouse NK-1.1 Antibody

|                           |  |
|---------------------------|--|
| <b>Catalog Number:</b>    | 203805, 203806                         |
| <b>Size:</b>              | 25 tests, 100 tests                    |
| <b>Target Name:</b>       | NK-1.1, CD161, NKR-P1C, NKR-P1B, Ly-55 |
| <b>Regulatory Status:</b> | RUO                                    |

### PRODUCT DETAILS

---


|                               |  |
|-------------------------------|--|
| <b>Clone:</b>                 | PK136  |
| <b>Application:</b>           | Flow Cytometry   |
| <b>Reactivity:</b>            | Mouse  |
| <b>Format:</b>                | FITC   |
| <b>Isotype:</b>               | Mouse IgG2a  |
| <b>Antibody Type:</b>         | Monoclonal   |
| <b>Formulation:</b>           | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA  |
| <b>Protein Concentration:</b> | Supplied at a lot-specific concentration.  |
| <b>Storage&amp;Handling:</b>  | The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.   |
| <b>Recommended Usage:</b>     | For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. FITC has an excitation max at 493 nm and an emission max at 525 nm. |
| <b>Excitation Laser:</b>      | Blue Laser (488 nm)  |
| <b>Isotype Control:</b>       | 301513   |

### BACKGROUND INFORMATION

---

NK-1.1, also known as CD161b/CD161c or Ly-55, is a type II transmembrane C-type lectin receptor encoded by the Klrb1 (NKR-P1) gene family in mice. It is predominantly expressed on natural killer (NK) cells and a subset of invariant NKT cells in selected mouse strains, including C57BL/6, FVB/N, and NZB. The activating isoform NKR-P1C lacks an inhibitory motif and promotes NK cell cytotoxicity, IFN- $\gamma$  production, and tumor cell lysis. NK-1.1 expression correlates with effective immune surveillance and graft rejection, making it a widely used marker for identifying and studying murine NK and NKT cell populations.

## PRODUCT DATA



Mouse splenocytes were stained with APC anti-Mouse CD49d and either FITC Anti-NK1.1 antibody clone PK136 (left panel) or an isotype control (right panel).