

## FITC Anti-Mouse/Human Ki-67 antibody

<b>Catalog Number:</b>	202411, 202412
<b>Size:</b>	25 tests, 100 tests
<b>Target Name:</b>	Ki-67, Ki67
<b>Regulatory Status:</b>	RUO

### PRODUCT DETAILS

---

<b>Clone:</b>	K67AR2b
<b>Application:</b>	Flow Cytometry
<b>Reactivity:</b>	Human, Mouse
<b>Format:</b>	FITC
<b>Isotype:</b>	Rat IgG2b
<b>Antibody Type:</b>	Monoclonal
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
<b>Protein Concentration:</b>	Supplied at a lot-specific concentration.
<b>Storage&amp;Handling:</b>	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
<b>Recommended Usage:</b>	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. FITC has an excitation max at 493 nm and an emission max at 525 nm.
<b>Excitation Laser:</b>	Blue Laser (488 nm)
<b>Isotype Control:</b>	303606

### BACKGROUND INFORMATION

---

Ki-67 is a nuclear protein strictly associated with cellular proliferation and is widely used as a biomarker in pathology and cancer research. It was first identified in the early 1980s during the search for antigens expressed in proliferating cells. The Ki-67 protein is encoded by the MKI67 gene located on chromosome 10 and is present during all active phases of the cell cycle (G1, S, G2, and M) but absent in resting (G0) cells. Because of this selective expression, Ki-67 serves as a reliable indicator of cell proliferation rates in both normal and tumor tissues.

Structurally, Ki-67 is a large protein, approximately 320–360 kDa, composed of multiple functional domains. It contains a forkhead-associated (FHA) domain and a series of tandem repeats known as the Ki-67 repeats, which are recognized by monoclonal antibodies used in immunohistochemistry. The protein is primarily localized within the nucleolus during interphase and associates with the perichromosomal layer during mitosis. Its exact molecular function remains under study, but evidence suggests that Ki-67 contributes to chromatin organization, particularly in maintaining the structural integrity of chromosomes during cell division.


Ki-67 does not have conventional extracellular ligands, as it functions within the nucleus. However, it interacts with chromatin and

several nuclear proteins that regulate the cell cycle and mitotic progression. Through these interactions, Ki-67 participates in organizing heterochromatin and possibly in ribosomal RNA synthesis regulation, facilitating efficient cell proliferation.

Clinically, Ki-67 has substantial value in oncology as a proliferation marker. High Ki-67 labeling indices (a measure of the percentage of Ki-67-positive tumor cells) often correlate with aggressive behavior and poor prognosis in diverse cancers, including breast, prostate, and neuroendocrine tumors. It is routinely used to assess tumor grade, guide therapeutic decisions, and monitor responses to treatment. Although not a direct therapeutic target, Ki-67's presence informs clinicians about tumor growth dynamics and potential responsiveness to cytotoxic therapies. Ongoing research explores whether inhibiting pathways regulating Ki-67 expression or function could complement conventional anticancer strategies, potentially yielding more selective antiproliferative therapies in the future.

## PRODUCT DATA

---



A-431 cells were fixed and permeabilized with 70% ethanol at -20°C for an hour and stained with FITC Anti-Mouse/Human Ki67 clone 11F6 (color-filled histogram) or isotype (gray histogram).

This product is supplied subject to the terms and conditions at [www.innocyto.com/web/terms.php](http://www.innocyto.com/web/terms.php) and may only be used as provided in the stated terms. Products are for Research Use Only.