

APC Anti-human HLA-G Antibody

Catalog Number:	112609, 112610
Size:	25 tests, 100 tests
Target Name:	HLA-G, Human Leukocyte Antigen-G
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	HLAGAM1
Application:	Flow Cytometry
Reactivity:	Human
Format:	APC
Isotype:	Mouse IgG1
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. APC has an excitation max at 650 nm and an emission max at 660 nm.
Excitation Laser:	Red Laser (633 nm)
Isotype Control:	301403

BACKGROUND INFORMATION

HLA-G is a non-classical major histocompatibility complex (MHC) class I molecule encoded within the human leukocyte antigen (HLA) region. Unlike classical MHC class I proteins, HLA-G has limited polymorphism and a restricted tissue distribution, primarily expressed at the maternal-fetal interface, where it plays a critical role in immune tolerance during pregnancy.

Structurally, HLA-G consists of a heavy chain associated with β 2-microglobulin and presents peptides similarly to other MHC class I molecules. However, alternative splicing generates multiple isoforms, including both membrane-bound (e.g., HLA-G1) and soluble forms (e.g., HLA-G5). These structural variants contribute to its diverse immunomodulatory functions.


HLA-G interacts with inhibitory receptors such as ILT2 (LILRB1), ILT4 (LILRB2), and KIR2DL4 expressed on immune cells including natural killer (NK) cells, T cells, and antigen-presenting cells. Through these ligand-receptor interactions, HLA-G suppresses immune responses by inhibiting cytotoxic activity, reducing cytokine production, and promoting regulatory cell phenotypes.

In disease contexts, aberrant expression of HLA-G has been associated with cancer, viral infections, and autoimmune disorders.

Many tumors exploit HLA-G expression to evade immune surveillance, leading to poorer clinical outcomes. Conversely, reduced HLA-G expression may contribute to pregnancy complications such as preeclampsia or recurrent miscarriage.

Therapeutically, HLA-G represents a promising target in both immunosuppression and immuno-oncology. Enhancing HLA-G activity could be beneficial in transplantation and autoimmune diseases by promoting immune tolerance. In contrast, blocking HLA-G or its receptors is being explored as a strategy to restore anti-tumor immunity. Ongoing research aims to better understand its mechanisms and develop targeted therapies that modulate HLA-G pathways.

PRODUCT DATA



JEG-3 cells were stained with APC Anti-Human HLA-G clone HLAGAM1 (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.