

PE Anti-Human CD40 Antibody

Catalog Number:	117003, 117004
Size:	25 tests, 100 tests
Target Name:	CD40, BP50, TNFRSF5
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	Dacetuzumab
Application:	Flow Cytometry
Reactivity:	Human
Format:	PE
Isotype:	Human IgG1
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage&Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. PE has an excitation max at 565 nm and an emission max at 575 nm.
Excitation Laser:	Blue Laser (488 nm) Green/Yellow laser (532/561nm)
Isotype Control:	301209

BACKGROUND INFORMATION

Human CD40 is a costimulatory receptor belonging to the tumor necrosis factor receptor (TNFR) superfamily, expressed on antigen-presenting cells such as B cells, dendritic cells, and macrophages, as well as some non-immune cells. It plays a central role in immune activation by promoting B-cell proliferation, immunoglobulin class switching, and dendritic cell maturation. Structurally, CD40 is a type I transmembrane protein with an extracellular domain composed of cysteine-rich repeats, a single transmembrane region, and a cytoplasmic tail that recruits adaptor proteins (e.g., TRAFs) to initiate downstream signaling pathways.

The primary ligand for CD40 is CD40L (CD154), expressed on activated T cells. Engagement of CD40 by CD40L triggers signaling cascades such as NF-κB activation, enhancing immune responses. Dysregulation of CD40 signaling is implicated in autoimmune diseases, chronic inflammation, and cancer. In oncology, CD40 can have dual roles: promoting anti-tumor immunity through immune activation, but also, in some contexts, supporting tumor cell survival.

Dacetuzumab is a humanized monoclonal antibody targeting CD40. It binds the extracellular domain and acts as a partial agonist, stimulating CD40 signaling while also mediating antibody-dependent cellular cytotoxicity (ADCC). Clinically, dacetuzumab has been

investigated in B-cell malignancies such as non-Hodgkin lymphoma. By activating immune cells and directly targeting CD40-expressing tumor cells, it represents an immunomodulatory therapeutic strategy, although its efficacy has been variable and is often explored in combination regimens.

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.