

Technical Data Sheet

Biotin Anti-Mouse/human CD45R/B220

Catalog Number: 200903, 200904

Size: 25 ug, 100 ug

Target Name: CD45R, B220

Regulatory Status: RUO

Product Details

Clone: RA3-6B2

Application: FC

Reactivity: Human, Mouse

Format: Biotin

Isotype: Rat IgG2a

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA

Protein Concentration: 0.5 mg/mL


Storage and Handling: The antibody solution should be stored between 2°C and 8°C

Recommended Usage: For flow cytometric staining, it is recommended to use less than 0.1 ug of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application.

Background Information

CD45R, also known as B220, is an isoform of CD45 (PTPRC), a member of the protein tyrosine phosphatase (PTP) family. With a molecular weight of approximately 180–240 kDa, CD45R is a transmembrane glycoprotein expressed on B cells at all developmental stages, as well as on activated B cells, subsets of T and NK cells, and certain abnormal T cell populations involved in systemic autoimmunity in MRL-Faslpr and MRL-Fasgld mice. Structurally, CD45R consists of an extracellular domain, a single transmembrane region, and two tandem intracytoplasmic catalytic domains. Functionally, CD45R plays a critical role in T cell receptor (TCR) and B cell receptor (BCR) signaling by regulating Src-family kinases, which are essential for lymphocyte activation and development. The primary ligands for CD45 include galectin-1, CD2, CD3, and CD4. CD45R is widely used as a pan-B cell marker, though CD19 may provide greater specificity for B cells.

Product Data



Mouse splenocytes stained with Biotin Anti-mouse/human CD45R_B220 clone RA3-6B2, followed by SA-PE.