www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

FITC Anti-Human CD32 Antibody

Catalog Number:	120603, 120604
Size:	25 tests, 100 tests
Target Name:	CD32, FcγRII, CD32a/FcγRIIa/FCGR2A, CD32b/FcγRIIb/FCGR2B
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	3D3
Application:	Flow Cytometry
Reactivity:	Human
Format:	FITC
Isotype:	Mouse IgG1
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. FITC has an excitation max at 493 nm and an emission max at 525 nm.
Excitation Laser:	Blue Laser (488 nm)
Isotype Control:	301415

BACKGROUND INFORMATION

Human CD32, also known as Fcγ receptor II (FcγRII), is a family of low-affinity receptors for the Fc portion of immunoglobulin G (IgG) that links antibody-mediated recognition to cellular immune responses. The family comprises three closely related receptors, CD32A (FcγRIIA), CD32B (FcγRIIB), and CD32C (FcγRIIC). They are expressed on various immune cells, including monocytes, macrophages, neutrophils, dendritic cells, and B cells, with expression patterns differing among the isoforms.


CD32 proteins are type I transmembrane glycoproteins belonging to the immunoglobulin superfamily. Their extracellular region contains two Ig-like domains that bind the Fc region of IgG, followed by a transmembrane segment and a cytoplasmic signaling region. CD32A and CD32C contain immunoreceptor tyrosine-based activation motifs (ITAMs), whereas CD32B contains an immunoreceptor tyrosine-based inhibitory motif (ITIM). CD32 has relatively low affinity for monomeric IgG but efficiently binds multivalent IgG-containing immune complexes.

The principal ligands are IgG antibodies and IgG-containing immune complexes, with receptor affinity varying according to IgG subclass and CD32 isoform. CD32A and CD32C generally promote cellular activation, phagocytosis, cytokine production, and

inflammatory responses, while CD32B counterbalances these activities and helps maintain immune tolerance.

Dysregulated CD32 signaling is implicated in autoimmune and inflammatory diseases, infectious disease, and cancer. In particular, CD32A genetic variants have been associated with susceptibility to several immune-mediated diseases, while CD32B contributes to regulation of B-cell and myeloid-cell activation. Therapeutically, CD32 is important both as a mediator of monoclonal-antibody activity and as a potential target itself. CD32B-directed antibodies have demonstrated antitumor activity in preclinical B-cell lymphoma models, while soluble FcγR-based approaches and CD32A antagonists are being investigated for antibody-driven autoimmune and inflammatory diseases.

PRODUCT DATA



Human Peripheral Blood Mononuclear Cells were stained with FITC Anti-Human CD32 clone 3D3 (color-filled histogram) or an isotype control (gray histogram). Histograms show CD32 expression in gated monocytes (left) and gated lymphocytes (right).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.