www.innocyto.com

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103
Irvine, CA 92618
Email: info@innocyto.com
Phone: 1-949-418-7543

APC/Cyanine7 Anti-human CD273 (B7-DC, PD-L2) Antibody

Catalog Number:	120809, 120810
Size:	25 tests, 100 tests
Target Name:	CD273, B7-DC, PD-L2, PDL2, B7DC
Regulatory Status:	RUO

PRODUCT DETAILS

Clone:	24F.10C12
Application:	Flow Cytometry
Reactivity:	Human
Format:	APC/Cyanine7
Isotype:	Mouse IgG2a
Antibody Type:	Monoclonal
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA
Protein Concentration:	Supplied at a lot-specific concentration.
Storage & Handling:	The antibody solution should be stored undiluted between 2°C and 8°C, and protected from prolonged exposure to light. Do not freeze.
Recommended Usage:	For flow cytometric staining, it is recommended to use 5 µL of this reagent per 0.5-1.0 million cells in a 100 µL volume. Optimal reagent performance should be determined by titration for each specific application. APC/Cyanine7 has an excitation max at 650 nm and an emission max at 774 nm.
Excitation Laser:	Red Laser (633 nm)
Isotype Control:	301518

BACKGROUND INFORMATION

Human programmed death-ligand 2 (PD-L2), also known as CD273 or B7-DC, is an immune checkpoint protein encoded by the PDCD1LG2 gene. It belongs to the B7 family of immunoregulatory molecules and plays an important role in regulating T-cell responses and maintaining immune balance. PD-L2 is expressed primarily by antigen-presenting cells, including dendritic cells and macrophages, and can also be induced on certain other cell types under inflammatory conditions. By interacting with its receptor, PD-L2 can modulate T-cell activation, cytokine production, and immune tolerance.


PD-L2 is a type I transmembrane glycoprotein composed of an extracellular region containing two immunoglobulin-like domains, a single transmembrane segment, and a short intracellular cytoplasmic tail. Its principal established receptor is programmed cell death protein 1 (PD-1, CD279), expressed on activated T cells and other immune cells. Binding of PD-L2 to PD-1 delivers inhibitory signals that reduce T-cell receptor signaling, cytokine production, and proliferation, thereby limiting excessive immune activation. PD-L2 has a higher binding affinity for PD-1 than its related ligand PD-L1 in many experimental systems. Additional interactions,

including binding to RGMb (repulsive guidance molecule b), have been reported and may contribute to immune regulation in particular tissues.

Dysregulated PD-L2 expression can influence disease progression. In cancer, some tumors and tumor-associated immune cells express PD-L2, enabling suppression of antitumor T-cell responses and potentially contributing to immune evasion. PD-L2 expression has been reported in several malignancies, although its prevalence and clinical significance vary by tumor type. In inflammatory and autoimmune conditions, PD-L2-mediated signaling may help restrain damaging immune responses, while altered regulation can affect immune tolerance and inflammation.

PD-L2 is a potential target for cancer immunotherapy. Blocking PD-L2, alone or alongside PD-1/PD-L1 pathway inhibitors, may restore antitumor T-cell activity in selected settings. Researchers are also investigating PD-L2 expression as a biomarker for immune activity and treatment response. Human PD-L2 is widely used in immunology and oncology research to study immune checkpoint signaling, T-cell regulation, and therapeutic strategies targeting the tumor immune microenvironment.

PRODUCT DATA



Human CD273 protein-coated beads (50%) mixed with non-coated beads (50%) were stained with APC/Cy7 Anti-Human CD273 clone 24F.10C12 (color-filled histogram) or an isotype control (gray histogram).

This product is supplied subject to the terms and conditions at www.innocyto.com/web/terms.php and may only be used as provided in the stated terms. Products are for Research Use Only.