

InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618

Technical Data Sheet

Biotin Anti-Mouse PD-L1

Catalog Number: 201303, 201304

Size: 25 ug, 100 ug

Target Name: CD274, PD-L1, B7-H1

Regulatory Status: RUO

Product Details

Clone: 10F.9G2
Application: FC
Reactivity: Mouse
Format: Biotin
Isotype: Rat IgG2b

Antibody Type: Monoclonal

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v)

BSA

Protein Concentration: 0.2 mg/mL

Storage and Handling: The antibody solution should be stored between 2°C and 8°C

Recommended Usage: For flow cytometric staining, it is recommended to use less than 0.1 ug of this reagent per 0.5-1.0 million cells in a 100 μ L volume. Optimal reagent performance should be

determined by titration for each specific application.

Background Information


PD-L1 (Programmed Death-Ligand 1), also known as CD274 or B7-H1, is a 40 kDa type I transmembrane protein belonging to the B7 family within the immunoglobulin receptor superfamily. The protein contains immunoglobulin V-like and C-like domains and is expressed by a wide range of hematopoietic and non-hematopoietic cells, including T cells, B cells, NK cells, dendritic cells, monocytes, endothelial cells, and various tumor cells. PD-L1 serves as a ligand for PD-1 (CD279) and plays a critical role in immune regulation by inhibiting T-cell activation, proliferation, and cytokine production upon engagement with PD-1. This interaction maintains immune homeostasis during infection or inflammation, preventing autoimmunity. However, in tumor microenvironments, PD-L1 expression enables immune evasion by suppressing cytotoxic T-cell function, contributing to tumor progression. PD-L1 expression is considered prognostic in several malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants. The PD-1/PD-L1 axis is a major target in cancer immunotherapy.

Product Data



InnoCyto Inc.

15375 Barranca Pkwy, Suite I-103 Irvine, CA 92618



Mouse splenocytes stained with either biotinylated anti-Mouse PD-L1 clone 10F.9G2 (blue histogram) or an isotype control (gray histogram), followed by SA-PE.