

## Technical Data Sheet

### Biotinylated Human Mesothelin (E296-G580) Protein (C-Fc-Avi)

**Catalog Number:** 801703, 801704  
**Size:** 25 ug, 100 ug  
**Target Name:** Mesothelin, MPF, MSLN  
**Regulatory Status:** RUO

#### Product Details

---

**Application:** ELISA, BLI  
**Format:** Liquid, Biotinylated  
**Expression Host:** HEK293  
**Species:** Human  
**Sources:** Human Mesothelin protein (Accession # AAH09272.1) (Glu296-Gly580) with C-terminus Fc-Avi tag is expressed in HEK293 cells. This protein was site-specifically labeled with Biotin by BirA ligase.  
**Accession Number:** Q13421  
**Molecular Weight:** The protein has a predicted molecular weight of 60kDa. Under DTT-reducing conditions, it migrates at approximately 65 kDa.  
**Affinity Tag:** C-Fc-Avi  
**Purity:** >95% based on SDS-PAGE under reducing condition  
**Formulation:** 1xPBS buffer, pH7.4, 0.22 µm filtered  
**Endotoxin level:** Not tested  
**Protein Concentration:** 25µg size is bottled at 0.2mg/mL concentration. 100 µg size is supplied at a lot-specific concentration.  
**Storage and Handling:** Briefly centrifuge the vial upon receipt. An unopened vial can be stored at 4°C for up to 2 weeks, or at -20°C or below for up to six months. The protein may be further diluted to 0.1 mg/mL using 0.22 µm-filtered PBS buffer (pH 7.4). For long-term storage, the diluted stock solution should be aliquoted and stored at ≤ -70°C to minimize freeze-thaw cycles. If additional dilution is required, carrier proteins such as FBS or BSA should be added to maintain protein stability.

#### Background Information

---

Mesothelin (MSLN), also known as CAK1, is a 70 kDa cell-surface protein cleaved into MPF (30 kDa) and mesothelin (40 kDa). MPF acts as a cytokine, while mesothelin is membrane-bound via glycosylphosphatidylinositol. It is overexpressed in various cancers, including mesotheliomas and ovarian cancer, and interacts with MUC16/CA-125, a cancer biomarker. Elevated mesothelin levels in serum make it a target for diagnostics and therapies, such as the recombinant immunotoxin SS1P, which has shown promise in clinical trials.

#### Product Data

---



Human Mesothelin (E296-G580, C-Fc-Avi) was biotinylated in vitro using BirA ligase. SDS-PAGE analysis under non-reducing (P-) conditions shows the protein has a purity greater than 95%. A gel shift assay using co-incubation with streptavidin indicates that the biotinylation efficiency of the Mesothelin protein exceeds 90%.